

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	CAC1B
<b>Synonyms</b>	BIII, CACNL1A5, CACNN, Cav2.2, DYT23, NEDNEH
<b>Description</b>	Human CAC1B full length protein-synthetic nanodisc
<b>Uniprot ID</b>	Q00975
<b>Protein Families</b>	Ion Channels: Calcium
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length CAC1B protein has a MW of 262.5kDa
<b>Delivery</b>	6~8weeks
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is the pore-forming subunit of an N-type voltage-dependent calcium channel, which controls neurotransmitter release from neurons. The encoded protein forms a complex with alpha-2, beta, and delta subunits to form the high-voltage activated channel. This channel is sensitive to omega-conotoxin-GVIA and omega-agatoxin-IIIa but insensitive to dihydropyridines. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2011]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

