

PRODUCT INFORMATION

CCL₅ **Target**

SISd; eoCP; SCYA5; RANTES; TCP228; D17S136E; **Synonyms**

SIS-delta

Recombinant human CCL5 Protein with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** P13501 **Expression Host HFK293**

C-Human Fc tag Tag

Molecular

CCL5(Ser24-Ser91) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of

34.0 kDa after removal of the signal peptide. The apparent molecular mass of CCL5-hFc is **Molecular Weight**

approximately 35-55 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue **Purity**

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before Formulation & lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and

Storage & Shipping

thawing). Lyophilized proteins are shipped at

ambient temperature.

This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member

of the CC subfamily, functions as a chemoattractant for blood monocytes, memory T helper cells and eosinophils. It causes the release **Background**

of histamine from basophils and activates eosinophils. This cytokine is one of the major HIVsuppressive factors produced by CD8+ cells. It functions as one of the natural ligands for the chemokine receptor chemokine (C-C motif) receptor 5 (CCR5), and it suppresses in vitro replication of the R5 strains of HIV-1, which use CCR5 as a coreceptor. Alternative splicing results

in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013]

Usage Research use only Conjugate Unconjugated

> Email: info@dimabio.com Website: www.dimabio.com










Figure 1. Human CCL5 Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

