

PRODUCT INFORMATION

| | |
|------------------------------|---|
| Tag | C-Flag Tag |
| Target | CCR3 |
| Synonyms | CC-CKR-3; C C CKR3; CD193; CKR 3; CKR3; CMKBR3 |
| Description | Human CCR3 full length protein-synthetic nanodisc |
| Delivery | In Stock |
| Uniprot ID | P51677 |
| Expression Host | HEK293 |
| Protein Families | Druggable Genome, GPCR, Transmembrane |
| Protein Pathways | Chemokine signaling pathway, Cytokine-cytokine receptor interaction |
| Molecular Weight | The human full length CCR3 protein has a MW of 41.0 kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein is a receptor for C-C type chemokines. It belongs to family 1 of the G protein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in eosinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. This receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cells in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other chemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alternatively spliced transcript variants have been described. |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure1. Elisa plates were pre-coated with Flag Tag CCR3-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR3-Nanodisc is 2.3ng/ml.



Figure2. Human CCR3-Nanodisc, Flag Tag on SDS-PAGE

