

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CCR6
Synonyms	BN-1; C-C CKR-6; CC-CKR-6; CCR-6; CD196; CKR-L3; CKRL3; CMKBR6; DCR2; DRY6; GPR29; GPRCY4; STRL22
Description	Human CCR6 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P51684
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
Molecular Weight	The human full length CCR6 protein has a MW of 42.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein is a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The gene is preferentially expressed by immature dendritic cells and memory T cells. The ligand of this receptor is macrophage inflammatory protein 3 alpha (MIP-3 alpha). This receptor has been shown to be important for B-lineage maturation and antigen-driven B-cell differentiation, and it may regulate the migration and recruitment of dendritic and T cells during inflammatory and immunological responses. Alternatively spliced transcript variants that encode the same protein have been described for this gene.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate CCR6-Nanodisc
0.2 μ g Human CCR6-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CCR6-Nanodisc (0.2 μ g/per well). Serial diluted anti-CCR6 monoclonal antibody (DMC100477) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-CCR6 monoclonal antibody binding with CCR6-Nanodisc is 2.955ng/ml.



Figure2. Human CCR6-Nanodisc, Flag Tag on SDS-PAGE

