

PRODUCT INFORMATION

CCR7 **Target**

BLR2; CC-CKR-7; CCR-7; CD197; CDw197; **Synonyms**

CMKBR7; EBI1

Human CCR7 full length protein-MNP **Description**

Delivery In Stock **Uniprot ID** P32248 **Expression Host HEK293**

Protein Families Druggable Genome, GPCR, Transmembrane

Chemokine signaling pathway, Cytokine-cytokine **Protein Pathways**

receptor interaction

The human full length CCR7 protein has a MW of **Molecular Weight**

42.9 kDa

Lyophilized from PBS. Normally 5% – 8% trehalose is added as protectants before Formulation &

Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Storage & Shipping

Lyophilized proteins are shipped at ambient

temperature.

The protein is a member of the G protein-coupled receptor family. This receptor was identified as a gene induced by the Epstein-Barr virus (EBV), and is thought to be a mediator of EBV effects on B lymphocytes. This receptor is expressed in various lymphoid tissues and activates B and T lymphocytes. It has been shown to control the migration of memory T cells to inflamed tissues, as well as stimulate dendritic cell maturation. The chemokine (C-C motif) ligand 19 (CCL19/ECL) has


been reported to be a specific ligand of this receptor. Signals mediated by this receptor regulate T cell homeostasis in lymph nodes, and

may also function in the activation and polarization of T cells, and in chronic

inflammation pathogenesis.

Usage Research use only

Background



Email: info@dimabio.com Website: www.dimabio.com





ELISA assay to evaluate CCR7-MNP 0.5µg Human CCR7-MNP per well




Figure 1. Elisa plates were pre-coated with 0.5µg/per well purified human CCR7 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CCR7 full length membrane nanoparticles is 5.185ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

