

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CCRL2
Synonyms	ACKR5, CKRX, CRAM, CRAM-A, CRAM-B, HCR
Description	Human CCRL2 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	O00421
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,Chemokines,Chemokine and Receptor,
Molecular Weight	The human full length CCRL2 protein has a MW of 39.5kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a chemokine receptor like protein, which is predicted to be a seven transmembrane protein and most closely related to CCR1. Chemokines and their receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. This gene is expressed at high levels in primary neutrophils and primary monocytes, and is further upregulated on neutrophil activation and during monocyte to macrophage differentiation. The function of this gene is unknown. This gene is mapped to the region where the chemokine receptor gene cluster is located. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

