Human CD147 Protein, hFc Tag Cat. No. PME100812



PRODUCT INFORMATION

| Target | CD147 |
|---------------------------------|---|
| Synonyms | Basigin;BSG;5F7;CD147;EMMPRIN;M6;OK;TCSF |
| Description | Recombinant Human CD147 Protein with C- terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P35613 |
| Expression Host | HEK293 |
| Тад | C-Human Fc Tag |
| Molecular Characterization | CD147(Gly140-Ala323) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 46.3 kDa after removal of the signal peptide.The apparent molecular mass of CD147-hFc is approximately 55-70 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene, basigin, is a plasma membrane protein that is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. Basigin is also a member of the immunoglobulin superfamily, ubiquitously expressed in various tissues. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2020] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| | |

Email: info@dimabio.com Website: www.dimabio.com



Human CD147 Protein, hFc Tag Cat. No. PME100812






Figure 1. Human CD147 Protein, hFc Tag on SDS-PAGE under reducing condition.

Email: info@dimabio.com Website: www.dimabio.com

