

PRODUCT INFORMATION

Target	CD164
Synonyms	LMOR;M-OR-1;MOP;MOR;MOR1;OPRM
Description	Recombinant Human CD164 with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q04900
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	CD164(Asp24-Asp162) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 40.7 kDa after removal of the signal peptide. The apparent molecular mass of CD164-hFc is approximately 70-130 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq, Oct 2013]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human CD164 Protein, hFc Tag on SDS-PAGE under reducing condition.

