Human CD44 Protein, His Tag Cat. No. PME100108



PRODUCT INFORMATION

Target	CD44
Synonyms	CD44;CDW44;CSPG8;ECMR-III;HCELL;HUTCH-I;IN;LHR;MC56;MDU2;MDU3;MIC4;Pgp1;Epican
Description	Recombinant Human CD44 Protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	P16070
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	CD44(Gln21-Pro220) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 22.9 kDa after removal of the signal peptide.The apparent molecular mass of CD44-His is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell
Background	interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis.
Usage	Research use only
Conjugate	Unconjugated



Figure 1. Human CD44 Protein, His Tag on SDS-PAGE under reducing condition. Address: Wuhan institute of Email: info@dimabio.com Biotechnology B7, Biolake No.666 Website: www.dimabio.com Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)





MARIO CONTRACTION

Email: info@dimabio.com Website: www.dimabio.com

