

**PRODUCT INFORMATION**

<b>Target</b>	CD45
<b>Synonyms</b>	B220;CD45;CD45R;GP180;L-CA;LCA;LY5;T200
<b>Description</b>	Recombinant Human CD45 with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P08575
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	CD45(Gln26-Lys577) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 61.6 kDa after removal of the signal peptide. The apparent molecular mass of CD45-His is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Background</b>	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human CD45 Protein, His Tag on SDS-PAGE under reducing condition.

### Human CD45, His Tagged protein ELISA

0.2  $\mu$ g of Human CD45, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CD45 Protein, His Tag(PME100581) can bind Anti-CD45 antibody(DM193), Rabbit mAb in a linear range of 0.64-16.00 ng/mL.

