

PRODUCT INFORMATION

CD93 **Target**

Synonyms C1qR(P);C1QR1;C1qRP;CDw93;dJ737E23.1;ECSM3;MXRA4

Recombinant human CD93 protein with C-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** Q9NPY3 **HEK293 Expression Host**

Tag C-Human Fc Tag

Molecular

Formulation &

Storage & Shipping

CD93(Thr22-Lys580) hFc(Glu99-Ala330) Characterization

Molecular Weight

The protein has a predicted molecular mass of 84.4 kDa after removal of the signal peptide. The apparent molecular mass of CD93-hFc is approximately 130-250

kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. **Purity**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions

Reconstitution of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form.

After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped

at ambient temperature.

The protein encoded by this gene is a cell-surface glycoprotein and type I membrane protein that was originally identified as a myeloid cell-specific marker. The encoded protein was once thought to be a receptor for C1q, but now is thought to instead be involved in intercellular adhesion and in the clearance of apoptotic cells. The intracellular cytoplasmic tail of this protein has

Background cells. The intracellular cytoplasmic tail of this protein has been found to interact with moesin, a protein known to

play a role in linking transmembrane proteins to the cytoskeleton and in the remodelling of the cytoskeleton.

> Email: info@dimabio.com Website: www.dimabio.com

[provided by RefSeq, Jul 2008]

Usage Research use only Conjugate Unconjugated








Figure 1. Human CD93 Protein, hFc Tag on SDS-PAGE under reducing condition

Human CD93, hFc Tagged protein ELISA

0.2 μg of Human CD93, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD93 Protein, hFc Tag (PME100689) can bind Anti-CD93(biosimilar) mAb (BME100176) in a linear range of 3.20–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

