

PRODUCT INFORMATION

CDH10 **Target Synonyms** CAD10

Recombinant Human CDH10 protein with C-Description

terminal 6×His tag

Delivery In Stock **Uniprot ID Q9Y6N8 Expression Host** HFK293 C-6×His Tag Tag

Molecular

Molecular Weight

CDH10(Gly55-Ala613) 6×His tag Characterization

The protein has a predicted molecular mass of

63.0 kDa after removal of the signal peptide. The apparent molecular mass of CDH10-His is

approximately 70-130 kDa due to glycosylation. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

Purity staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a type II classical cadherin of the cadherin superfamily. Alternative splicing of this gene results in multiple transcript variants. At

least one of these variants encodes a preproprotein that is proteolytically processed to generate the mature cadherin protein. These integral membrane proteins mediate calciumdependent cell-cell adhesion and are composed of a large N-terminal extracellular domain, a single membrane-spanning domain, and a small, highly conserved C-terminal cytoplasmic domain. The extracellular domain consists of 5 subdomains,

Background each containing a cadherin motif, and appears to

determine the specificity of the protein's homophilic cell adhesion activity. Type II (atypical) cadherins are defined based on their lack of a histidine-alanine-valine (HAV) cell adhesion recognition sequence specific to type I

cadherins. This particular cadherin is predominantly expressed in brain and is putatively involved in synaptic adhesions, axon outgrowth and guidance. Mutations in this gene may be associated with lung squamous cell carcinoma and colorectal cancer in human patients. [provided by RefSeq, Nov 2015]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Human CDH10 Protein, His Tag on SDS-PAGE under reducing condition.



