

PRODUCT INFORMATION

| | |
|---|--|
| Target | CDH17 |
| Synonyms | HPT-1; HPT1 |
| Description | Recombinant human CDH17(668-777) Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | Q12864 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | CDH17(Ala668-Gly777) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 38.3 kDa after removal of the signal peptide. The apparent molecular mass of CDH17(668-777)-hFc is approximately 35-55 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. The encoded protein is cadherin-like, consisting of an extracellular region, containing 7 cadherin domains, and a transmembrane region but lacking the conserved cytoplasmic domain. The protein is a component of the gastrointestinal tract and pancreatic ducts, acting as an intestinal proton-dependent peptide transporter in the first step in oral absorption of many medically important peptide-based drugs. The protein may also play a role in the morphological organization of liver and intestine. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human CDH17(668-777) Protein, hFc Tag on SDS-PAGE under reducing condition.

