

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	CHRM2
Synonyms	HM2
Description	Human CHRM2 full length protein-synthetic nanodisc
Uniprot ID	P08172
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Calcium signaling pathway, Neuroactive ligand-receptor interaction, Regulation of actin cytoskeleton
Molecular Weight	The human full length CHRM2 protein has a MW of 51.7 kDa
Delivery	3-4 weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine to these receptors and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarinic cholinergic receptor 2 is involved in mediation of bradycardia and a decrease in cardiac contractility. Multiple alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate CHRM2-Nanodisc 0.2 μ g Human CHRM2-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CHRM2-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CHRM2-Nanodisc is 13.52ng/ml.

kDa M R



Figure2. Human CHRM2-Nanodisc, Flag Tag on SDS-PAGE

