

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	CLCN3
Synonyms	CLC3, CIC-3
Description	Human CLCN3-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P51790
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length CLCN3-Strep protein has a MW of 91 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the voltage-gated chloride channel (CIC) family. The encoded protein is present in all cell types and localized in plasma membranes and in intracellular vesicles. It is a multi-pass membrane protein which contains a CIC domain and two additional C-terminal CBS (cystathionine beta-synthase) domains. The CIC domain catalyzes the selective flow of Cl ⁻ ions across cell membranes, and the CBS domain may have a regulatory function. This protein plays a role in both acidification and transmitter loading of GABAergic synaptic vesicles, and in smooth muscle cell activation and neointima formation. This protein is required for lysophosphatidic acid (LPA)-activated Cl ⁻ current activity and fibroblast-to-myofibroblast differentiation. The protein activity is regulated by Ca ²⁺ /calmodulin-dependent protein kinase II (CaMKII) in glioma cells. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Aug 2011]
Usage	Research use only
Conjugate	Unconjugated

