

PRODUCT INFORMATION

CLDN4 **Target**

Synonyms CPE-R; CPETR; CPETR1; hCPE-R; WBSCR8 Recombinant Human CLDN4(145-157) Protein

with C-terminal human Fc tag

Delivery In Stock **Uniprot ID** 014493 **Expression Host HEK293**

Tag C-Human Fc Tag

Molecular

Background

Description

CLDN4(Gln145-Lys157) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of **Molecular Weight**

27.6 kDa after removal of the signal peptide. The apparent molecular mass of CLDN4(145-157)-hFc is approximately 25-35 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & Reconstitution

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this intronless gene belongs to the claudin family. Claudins are


integral membrane proteins that are components of the epithelial cell tight junctions, which regulate movement of solutes and ions through the paracellular space. This protein is a high-affinity receptor for Clostridium perfringens enterotoxin (CPE) and may play a role in internal

organ development and function during pre- and postnatal life. This gene is deleted in Williams-Beuren syndrome, a neurodevelopmental

disorder affecting multiple systems. [provided by RefSeq, Sep 2013]

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only



/+86-400-006-0995(China)






Figure 1. Human CLDN4(145-157) Protein, hFc Tag on SDS-PAGE under reducing condition.



