

## **PRODUCT INFORMATION**

CLEC4C **Target** 

**Synonyms** BDCA-2;CD303

Recombinant human CLEC4C protein with N-**Description** 

terminal human Fc tag

**Delivery** In Stock **Uniprot ID** Q8WTT0 **Expression Host HEK293** 

Tag N-Human Fc Tag

Molecular

**Molecular Weight** 

hFc(Glu99-Ala330) CLEC4C(Asn45-Ile213) Characterization

The protein has a predicted molecular mass of 46.1 kDa after removal of the signal peptide. The

apparent molecular mass of hFc-CLEC4C is approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a

common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling,


glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 **Background** transmembrane protein may play a role in

dendritic cell function. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]

> Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Unconjugated Conjugate



/+86-400-006-0995(China)









Figure 1. Human CLEC4C Protein, hFc Tag on SDS-PAGE under reducing condition.

## Human CLEC4C, hFc Tagged protein ELISA

0.2 μg of Human CLEC4C, hFc tagged protein per well



Anti-CLEC4C (litifilimab biosimilar) mAb (ng/mL)

Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CLEC4C Protein, hFc Tag (PME100756) can bind Anti-CLEC4C (litifilimab biosimilar) mAb (BME100155) in a linear range of 0.64–80.00 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

