

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	CLIC2
Synonyms	CLCNL2, CLIC2b, MRXS32, XAP121
Description	Human CLIC2 full length protein-synthetic nanodisc
Uniprot ID	O15247
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length CLIC2 protein has a MW of 28.4kDa
Delivery	6~8weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a chloride intracellular channel protein. Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. This protein plays a role in inhibiting the function of ryanodine receptor 2. A mutation in this gene is the cause of an X-linked form of cognitive disability. [provided by RefSeq, Jul 2017]
Usage	Research use only
Conjugate	Unconjugated

