

PRODUCT INFORMATION

| Тад | C-Flag Tag |
|---------------------------------|---|
| Target | СМТМ6 |
| Synonyms | CKLFSF6; PRO2219 |
| Description | Human CMTM6 full length protein-synthetic nanodisc |
| Delivery | 3-4 weeks |
| Uniprot ID | Q9NX76 |
| Expression Host | HEK293 |
| Protein Families | Transmembrane |
| Protein Pathways | N/A |
| Molecular Weight | The human full length CMTM6 protein has a MW |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in |
| Storage & Shipping | lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene belongs to the chemokine-like factor gene superfamily, a novel family that is similar to the chemokine and transmembrane 4 superfamilies. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 3. This gene is widely expressed in many tissues, but the exact function of the encoded protein is unknown. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |
| | |










Figure1. Elisa plates were pre-coated with Flag Tag CMTM6-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CMTM6-Nanodisc is 4.393ng/ml.




Figure2. Human CMTM6-Nanodisc, Flag Tag on SDS-PAGE

Email: info@dimabio.com Website: www.dimabio.com

