

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CX3CR1
Synonyms	CCRL1; CMKBRL1; CMKDR1; GPR13; GPRV28; V28
Description	Human CX3CR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P49238
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Chemokine signaling pathway, Cytokine-cytokine receptor interaction
Molecular Weight	The human full length CX3CR1 protein has a MW of 40.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Fractalkine is a transmembrane protein and chemokine involved in the adhesion and migration of leukocytes. The protein encoded by this gene is a receptor for fractalkine. The encoded protein also is a coreceptor for HIV-1, and some variations in this gene lead to increased susceptibility to HIV-1 infection and rapid progression to AIDS.
Usage	Research use only
Conjugate	Unconjugated



**ELISA assay to evaluate CX3CR1-Nanodisc**  
0.2µg Human CX3CR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag CX3CR1-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with CX3CR1-Nanodisc is 2.796ng/ml.



Figure2. Human CX3CR1-Nanodisc, Flag Tag on SDS-PAGE

