

PRODUCT INFORMATION

| | |
|------------------------------|--|
| Target | CXADR |
| Synonyms | CAR;CAR4/6;HCAR |
| Description | Recombinant Human CXADR Protein with C-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | P78310 |
| Expression Host | HEK293 |
| Tag | C-6×His Tag |
| Molecular Characterization | CXADR(Leu20-Gly237) 6×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 24.9 kDa after removal of the signal peptide. The apparent molecular mass of CXADR-His is approximately 25-35 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | The protein encoded by this gene is a type I membrane receptor for group B coxsackieviruses and subgroup C adenoviruses. Several transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene are found on chromosomes 15, 18, and 21. [provided by RefSeq, May 2011] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1.Human CXADR Protein, His Tag on SDS-PAGE under reducing condition.

