

**PRODUCT INFORMATION**

<b>Target</b>	CXCL3
<b>Synonyms</b>	C-X-C Motif Chemokine 3;GRO-Gamma (1-73);Growth-Regulated Protein Gamma;GRO-Gamma;Macrophage Inflammatory Protein 2-Beta;MIP2-Beta;GRO-Gamma (5-73);CXCL3;GRO3;GROG;SCYB3
<b>Description</b>	Recombinant Human C-X-C Motif Chemokine 3 is produced by our E.coli expression system and the target gene encoding Ala35-Asn107 is expressed with a 6His tag at the N-terminus.
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P19876
<b>Expression Host</b>	E.coli
<b>Tag</b>	
<b>Molecular Characterization</b>	Not available
<b>Molecular Weight</b>	10.1 KDa
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Background</b>	C-X-C Motif Chemokine 3 (CXCL3) is a secreted protein that belongs to the intercrine alpha (chemokine CXC) family. CXCL3 controls the migration and adhesion of monocytes and mediates its effect on its target cell by interacting with a cell surface chemokine receptor called CXCR2. In addition, CXCL3 is thought to play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Greater than 95% as determined by reducing SDS-PAGE.

DIMABIO CONFIDENTIAL

