

## **PRODUCT INFORMATION**

CXCR4 Target

Synonyms CD184;D2S201E;FB22;HM89;HSY3RR;LAP-3;LAP3;LCR1;LESTR;NPY3R;NPYR;NPYRL;NPYY3R;WHIM;WHIMS

Recombinant Human CXCR4 with C-terminal human Fc tag Description

Delivery **Uniprot ID** P61073 **Expression Host** HEK293 Tag C-Human Fc Tag

Molecular Characterization CXCR4(Met1-Lvs38) hFc(Glu99-Ala330)

The protein has a predicted molecular mass of 30.5 kDa after removal of the signal peptide. The apparent molecular mass of CXCR4-hFc is approximately 35-55 kDa due to glycosylation. **Molecular Weight** 

Purity The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a CXC chemokine receptor specific for stromal cell-derived factor-1. The protein has 7 transmembrane regions and is located on the cell surface. It acts with the CD4 protein to support HIV entry into cells and is also highly expressed in breast cancer cells. Mutations in this gene have been associated with WHIM (warts, hypogammaglobulinemia, infections, and myelokathexis) syndrome. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]

Usage Research use only

Background

Conjugate Unconjugated




Figure 1. Human CXCR4 Protein, hFc Tag on SDS-PAGE under reducing condition.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

