

PRODUCT INFORMATION

Target	CYSLTR1
Synonyms	CYSLT1, CYSLTR, CYSLT1R, HMTMF81
Description	Recombinant human CYSLTR1 Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9Y271
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	CYSLTR1(Met1-Ser27) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 29.2 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	This gene encodes a member of the G-protein coupled receptor 1 family. The encoded protein is a receptor for cysteinyl leukotrienes, and is involved in mediating bronchoconstriction via activation of a phosphatidylinositol-calcium second messenger system. Activation of the encoded receptor results in contraction and proliferation of bronchial smooth muscle cells, eosinophil migration, and damage to the mucus layer in the lung. Upregulation of this gene is associated with asthma and dysregulation may also be implicated in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2013]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human CYSLTR1 Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

