

## **PRODUCT INFORMATION**

DKK1 **Target** 

**Synonyms** DKK-1;SK

Recombinant human DKK1 Protein with C-**Description** 

terminal human Fc tag

**Delivery** In Stock **Uniprot ID** 094907 **Expression Host HEK293** 

Tag C-Human Fc Tag

Molecular

Storage & Shipping

**Background** 

DKK1(Thr32-His266) hFc(Glu99-Ala330) Characterization

The protein has a predicted molecular mass of

51.9 kDa after removal of the signal peptide. The apparent molecular mass of DKK1-hFC is **Molecular Weight** 

approximately 70 kDa due to glycosylation. The purity of the protein is greater than 90% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the dickkopf family of proteins. Members of this family are secreted proteins characterized by two cysteinerich domains that mediate protein-protein interactions. The encoded protein binds to the LRP6 co-receptor and inhibits beta-catenin-dependent Wnt signaling. This gene plays a role

in embryonic development and may be important in bone formation in adults. Elevated expression of this gene has been observed in numerous human cancers and this protein may promote proliferation, invasion and growth in cancer cell lines.

> Email: info@dimabio.com Website: www.dimabio.com

**Usage** Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)








Figure 1. Human DKK1 Protein, hFc Tag on SDS-PAGE under reducing condition.

## Human DKK1, hFc Tagged protein ELISA

0.2 μg of Human LRP6, His tagged protein per well




Figure 2. ELISA plate pre-coated by 5  $\mu$ g/mL (100  $\mu$ L/well) Human LRP6 Protein, His Tag (PME101193) can bind Human DKK1 Protein, hFc Tag(PME100230) in a linear range of 16–400ng/mL.



66 nina Email: info@dimabio.com Website: www.dimabio.com



## Human DKK1, hFc Tagged protein ELISA

 $0.2~\mu g$  of Human LRP5(644-1263), His tagged protein per well




Figure 3. ELISA plate pre-coated by 5  $\mu$ g/mL (100  $\mu$ L/well) Human LRP5(644-1263)Protein, His Tag can bind Human DKK1 Protein, HFC Tag(PME100230) in a linear range of 16–400ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

