

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	DRD1
Synonyms	DADR; DRD1A
Description	Human DRD1-Strep full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P21728
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Calcium signaling pathway, Gap junction, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length DRD1-Strep protein has a MW of 49.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This protein is D1 subtype of the dopamine receptor. The D1 subtype is the most abundant dopamine receptor in the central nervous system. This G-protein coupled receptor stimulates adenylyl cyclase and activates cyclic AMP-dependent protein kinases. D1 receptors regulate neuronal growth and development, mediate some behavioral responses, and modulate dopamine receptor D2-mediated events.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate DRD1-Strep-Nanodisc
0.2µg Human DRD1-Strep-Nanodisc per well




Figure 1. Elisa plates were pre-coated with C-Flag&Strep Tag DRD1-Strep-Nanodisc (0.2µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with DRD1-Strep-nanodisc is 3.600ng/ml.




Figure 2. Human DRD1-Strep-Nanodisc, C-Flag&Strep Tag on SDS-PAGE

