

PRODUCT INFORMATION

Target	ECSCR
Synonyms	ARIA;ECSM2
Description	Recombinant Human ECSCR Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q19T08
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	ECSCR(Gln25-Val122) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 36.0 kDa after removal of the signal peptide. The apparent molecular mass of ECSCR-hFc is approximately 35-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is primarily found in endothelial cells and blood vessels, where it is involved in cell shape changes and EGF-induced cell migration. It can enhance the activation of vascular endothelial growth factor receptor-2/kinase insert domain receptor and also promote the proteolysis of internalized kinase insert domain receptor. This gene may play a role in angiogenesis-related diseases. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human ECSCR Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

