

PRODUCT INFORMATION

EDA Target

Synonyms ED1;EDA2

Recombinant Human EDA Protein with N-terminal **Description**

human Fc tag

Delivery In Stock **Uniprot ID** Q92838 **Expression Host HEK293**

Tag N-Human Fc Tag

Molecular

Molecular Weight

Background

hFc(Glu99-Ala330) EDA(Ser160-Ser391) Characterization

The protein has a predicted molecular mass of 50.3 kDa after removal of the signal peptide. The

apparent molecular mass of hFc-EDA is

approximately 55-70 kDa due to glycosylation. The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

Purity

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a type II membrane protein that can be cleaved by furin to produce a secreted form. The encoded protein, which belongs to the tumor necrosis factor family, acts as a homotrimer and may be involved in cell-

cell signaling during the development of ectodermal organs. Defects in this gene are a

cause of ectodermal dysplasia, anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this

> Email: info@dimabio.com Website: www.dimabio.com

gene. [provided by RefSeq, Jul 2008]

Research use only Usage

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)








Figure 1.Human EDA Protein, hFc Tag on SDS-PAGE under reducing condition.

Human EDA, hFc Tagged protein ELISA

0.2 μg of Human EDA, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human EDA Protein, hFc Tag (PME100661) can bind Anti-EDA antibody(DMC388), IgG1 Chimeric mAb in a linear range of 3.20–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

