

PRODUCT INFORMATION

Target	EDA
Synonyms	ED1;EDA2
Description	Recombinant Human EDA Protein with N-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q92838
Expression Host	HEK293
Tag	N-Human Fc Tag
Molecular Characterization	hFc(Glu99-Ala330) EDA(Ser160-Ser391)
Molecular Weight	The protein has a predicted molecular mass of 50.3 kDa after removal of the signal peptide. The apparent molecular mass of hFc-EDA is approximately 55-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	The protein encoded by this gene is a type II membrane protein that can be cleaved by furin to produce a secreted form. The encoded protein, which belongs to the tumor necrosis factor family, acts as a homotrimer and may be involved in cell-cell signaling during the development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia, anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human EDA Protein, hFc Tag on SDS-PAGE under reducing condition.

Human EDA, hFc Tagged protein ELISA

0.2 µg of Human EDA, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human EDA Protein, hFc Tag (PME100661) can bind Anti-EDA antibody(DMC388), IgG1 Chimeric mAb in a linear range of 3.20-80 ng/mL.

