

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	EDNRB
Synonyms	ABCDS, ET-B, ET-BR, ETB, ETB1, ETBR, ETRB, HSCR, HSCR2, WS4A
Description	Human EDNRB-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P24530
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome, GPCRDB Class A Rhodopsin-like,Peptide
Protein Pathways	GPCRs,Prostaglandin synthesis regulation,Angiogenesis,Cancer,Endothelial Cell Biology,
Molecular Weight	The human full length EDNRB-Strep protein has a MW of 49.6 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	The protein encoded by this gene is a G protein- coupled receptor which activates a phosphatidylinositol-calcium second messenger system. Its ligand, endothelin, consists of a family of three potent vasoactive peptides: ET1, ET2, and ET3. Studies suggest that the multigenic disorder, Hirschsprung disease type 2, is due to mutations in the endothelin receptor type B gene. Alternative splicing and the use of alternative promoters results in multiple transcript variants. [provided by RefSeq, Oct 2016]
Usage	Research use only
Conjugate	Unconjugated

