

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	EMP2
Synonyms	XMP
Description	Human EMP2 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P54851
Expression Host	HEK293
Protein Families	Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length EMP2 protein has a MW of 19.2 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Formulation & Reconstitution	A tetraspan protein of the PMP22/EMP family. The encoded protein regulates cell membrane composition. It has been associated with various functions including endocytosis, cell signaling, cell proliferation, cell migration, cell adhesion, cell death, cholesterol homeostasis, urinary albumin excretion, and embryo implantation. It is known to negatively regulate caveolin-1, a scaffolding protein which is the main component of the caveolae plasma membrane invaginations found in most cell types. Through activation of PTK2 it positively regulates vascular endothelial growth factor A. It also modulates the function of specific integrin isomers in the plasma membrane. Up-regulation of this gene has been linked to cancer progression in multiple different tissues. Mutations in this gene have been associated with nephrotic syndrome type 10 (NPHS10). Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	
Storage&Shipping	
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate EMP2-Nanodisc
0.2 μ g Human EMP2-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag EMP2-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with EMP2-Nanodisc is 4.249ng/ml.



Figure2. Human EMP2-Nanodisc, Flag Tag on SDS-PAGE

