

PRODUCT INFORMATION

Target	EPHA3
Synonyms	EK4; ETK; HEK; ETK1; HEK4; TYRO4
Description	Recombinant human EPHA3(436-541) Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P29320
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	EPHA3(Ala436-Gln541) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 38.1 kDa after removal of the signal peptide. The apparent molecular mass of EPHA3(436-541)-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. This gene encodes a protein that binds ephrin-A ligands. Two alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human EPHA3(436-541) Protein, hFc Tag on SDS-PAGE under reducing condition.

