

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag Tag
<b>Target</b>	F2RL3
<b>Synonyms</b>	PAR4
<b>Description</b>	Human F2RL3 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q96RI0
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
<b>Protein Pathways</b>	Neuroactive ligand-receptor interaction
<b>Molecular Weight</b>	The human full length F2RL3 protein has a MW of 41.1 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	A member of the protease-activated receptor subfamily, part of the G-protein coupled receptor 1 family of proteins. The encoded receptor is proteolytically processed to reveal an extracellular N-terminal tethered ligand that binds to and activates the receptor. This receptor plays a role in blood coagulation, inflammation and response to pain. Hypomethylation at this gene may be associated with lung cancer in human patients.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



**ELISA assay to evaluate F2RL3-Nanodisc**  
0.2 $\mu$ g Human F2RL3-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag F2RL3-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with F2RL3-Nanodisc is 3.115ng/ml.




Figure2. Human F2RL3-Nanodisc, Flag Tag on SDS-PAGE

