

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	FFAR1
Synonyms	FFA1R; GPCR40; GPR40
Description	Human FFAR1 full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	O14842
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length FFAR1 protein has a MW of 31.5 kDa
Background	A member of the GP40 family of G protein-coupled receptors that are clustered together on chromosome 19. The encoded protein is a receptor for medium and long chain free fatty acids and may be involved in the metabolic regulation of insulin secretion. Polymorphisms in this gene may be associated with type 2 diabetes.
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate FFAR1-Nanodisc 0.2 μ g Human FFAR1-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag FFAR1-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FFAR1-Nanodisc is 4.566ng/ml.



Figure2. Human FFAR1-Nanodisc, Flag Tag on SDS-PAGE

