

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	FFAR4
<b>Synonyms</b>	BMIQ10; GPR120; GPR129; GT01; O3FAR1; PGR4
<b>Description</b>	Human FFAR4 full length protein-synthetic nanodisc
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q5NUL3
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, Transmembrane
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length FFAR4 protein has a MW of 42.2 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein is a G protein-coupled receptor (GPR) which belongs to the rhodopsin family of GPRs. The encoded protein functions as a receptor for free fatty acids, including omega-3, and participates in suppressing anti-inflammatory responses and insulin sensitizing.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated



### ELISA assay to evaluate FFAR4-Nanodisc 0.2 $\mu$ g Human FFAR4-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag FFAR4-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FFAR4-Nanodisc is 20.10ng/ml.




Figure 2. Human FFAR4-Nanodisc, Flag Tag on SDS-PAGE

