

PRODUCT INFORMATION

Target	FGFR2
Synonyms	FGFR2IIIb; BEK; JWS; BBDS; CEK3; CFD1; ECT1; KGFR; TK14; TK25; BFR-1; CD332; K-SAM
Description	Recombinant human FGFR2b Protein with C-terminal 10×His tag
Delivery	In Stock
Uniprot ID	P21802-3
Expression Host	HEK293
Tag	C-10×His tag
Molecular Characterization	FGFR2b(Arg22-Asp375) 10×His tag
Molecular Weight	The protein has a predicted molecular mass of 40.6 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Jan 2009]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human FGFR2b Protein, His Tag on SDS-PAGE under reducing condition.

Human FGFR2, His Tagged protein ELISA

0.2 μg of Human FGFR2, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2 $\mu\text{g}/\text{mL}$ (100 $\mu\text{L}/\text{well}$) Human FGFR2b Protein, His Tag (PME101408) can bind Anti-FGFR2(bemarituzumab biosimilar) mAb (BME100260) in a linear range of 0.13-80 ng/mL.

