

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	FSHR
Synonyms	FSHR1; FSHRO; LGR1; ODG1
Description	Human FSHR full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P23945
Expression Host	HEK293
Protein Families	Druggable Genome, ES Cell Differentiation/IPS, GPCR, Transmembrane
Protein Pathways	Neuroactive ligand-receptor interaction
Molecular Weight	The human full length FSHR protein has a MW of 78.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	
Background	The protein belongs to family 1 of G-protein coupled receptors. It is the receptor for follicle stimulating hormone and functions in gonad development. Mutations in this gene cause ovarian dysgenesis type 1, and also ovarian hyperstimulation syndrome. Alternative splicing results in multiple transcript variants.
Usage	Research use only
Conjugate	Unconjugated



ELISA assay to evaluate FSHR-Nanodisc 0.2 μ g Human FSHR-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag FSHR-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC₅₀ for anti-Flag monoclonal antibody binding with FSHR-Nanodisc is 3.687ng/ml.



Figure2. Human FSHR-Nanodisc, Flag Tag on SDS-PAGE

