

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	FXYD3
<b>Synonyms</b>	MAT8, PLML
<b>Description</b>	Human FXYD3 full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	Q14802
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Other
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length FXYD3 protein has a MW of 9.3kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene belongs to a small family of FXYD-domain containing regulators of Na <sup>+</sup> /K <sup>+</sup> ATPases which share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD, and containing 7 invariant and 6 highly conserved amino acids. This gene encodes a cell membrane protein that may regulate the function of ion-pumps and ion-channels. This gene may also play a role in tumor progression. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Oct 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

