

PRODUCT INFORMATION

Target	FZD10
Synonyms	CD350;FZ-10;Fz10;FzE7;hFz10
Description	Recombinant Human FZD10 with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9ULW2
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	FZD10(Ile21-Gly161) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 42.2 kDa after removal of the signal peptide. The apparent molecular mass of FZD10-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene is a member of the frizzled gene family. Members of this family encode 7-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. Using array analysis, expression of this intronless gene is significantly up-regulated in two cases of primary colon cancer. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human FZD10 Protein, hFc Tag on SDS-PAGE under reducing condition.

**Human FZD10, hFc Tagged protein ELISA**

0.2 µg of Human FZD10, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human FZD10 Protein, hFc Tag (PME100650) can bind Anti-FZD10(tabituximab biosimilar) mAb (BME100185) in a linear range of 3.20-80 ng/mL.

