

PRODUCT INFORMATION

Tag	C-Flag Tag
Expression Host	HEK293
Target	FZD3
Synonyms	Fz-3
Description	Human FZD3 full length protein-synthetic nanodisc
Uniprot ID	Q9NPG1
Protein Families	GPCR, Transmembrane, Druggable Genome,
Protein Pathways	Wnt signaling, Wnt signaling and pluripotency, Cancer, Notch, Wnt Pathway, Stem Cell ,
Molecular Weight	The human full length FZD3 protein has a MW of 76.3kDa
Delivery	6~8weeks
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 μ m) prior to use.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene is a member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. The function of this protein is unknown, although it may play a role in mammalian hair follicle development. Alternative splicing results in multiple transcript variants. This gene is a susceptibility locus for schizophrenia. [provided by RefSeq, Dec 2010]
Usage	Research use only
Conjugate	Unconjugated

