

PRODUCT INFORMATION

Target	FZD4
Synonyms	CD344; EVR1; FEVR; Fz-4; Fz4; FZD4S; FzE4; GPCR; hFz4
Description	Human FZD4 full length protein-MNP
Delivery	In Stock
Uniprot ID	Q9ULV1
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Basal cell carcinoma, Colorectal cancer, Melanogenesis, Pathways in cancer, Wnt signaling pathway
Molecular Weight	The human full length FZD4 protein has a MW of 60.3 kDa
Formulation & Reconstitution	Lyophilized from PBS. Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	A member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. This protein may play a role as a positive regulator of the Wingless type MMTV integration site signaling pathway. A transcript variant retaining intronic sequence and encoding a shorter isoform has been described, however, its expression is not supported by other experimental evidence.
Usage	Research use only
Conjugate	Unconjugated



**ELISA assay to evaluate FZD4-MNP**  
0.5µg Human FZD4-MNP per well




Figure1. Elisa plates were pre-coated with 0.5µg/per well purified human FZD4 full length membrane nanoparticles. Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with FZD4 full length membrane nanoparticles is 13.75ng/ml.

