

## PRODUCT INFORMATION

<b>Tag</b>	C-Flag&Strep Tag
<b>Target</b>	FZD9
<b>Synonyms</b>	CD349, FZD3
<b>Description</b>	Human FZD9-Strep full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	O00144
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	GPCR,Transmembrane,Druggable Genome, Wnt NetPath 8,Wnt signaling,Wnt signaling and pluripotency,Cancer,Notch,Wnt Pathway,Stem Cell ,
<b>Protein Pathways</b>	
<b>Molecular Weight</b>	The human full length FZD9-Strep protein has a MW of 64.5 kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD9 gene is located within the Williams syndrome common deletion region of chromosome 7, and heterozygous deletion of the FZD9 gene may contribute to the Williams syndrome phenotype. FZD9 is expressed predominantly in brain, testis, eye, skeletal muscle, and kidney. [provided by RefSeq, Jul 2008]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

