

PRODUCT INFORMATION

| | |
|---|--|
| Target | GAL |
| Synonyms | ETL8; GALN; GLNN; GMAP; GAL-GMAP |
| Description | Recombinant human GAL Protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P22466 |
| Expression Host | HEK293 |
| Tag | C-Human Fc tag |
| Molecular Characterization | GAL(Ala20-Ser123) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 37.6 kDa after removal of the signal peptide. The apparent molecular mass of GAL-hFc is approximately 35-55 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a neuroendocrine peptide that is widely expressed in the central and peripheral nervous systems and also the gastrointestinal tract, pancreas, adrenal gland and urogenital tract. The encoded protein is a precursor that is proteolytically processed to generate two mature peptides: galanin and galanin message-associated peptide (GMAP). Galanin has diverse physiological functions including nociception, feeding and energy homeostasis, osmotic regulation and water balance. GMAP has been demonstrated to possess antifungal activity and hypothesized to be part of the innate immune system. [provided by RefSeq, Jul 2015] |
| Usage | Research use only |
| Conjugate | Unconjugated |





Figure 1. Human GAL Protein, hFc Tag on SDS-PAGE under reducing condition.

