

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag&Strep Tag
<b>Expression Host</b>	HEK293
<b>Target</b>	GIPR
<b>Synonyms</b>	PGQTL2
<b>Description</b>	Human GIPR-Strep full length protein-synthetic nanodisc
<b>Uniprot ID</b>	P48546
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
<b>Protein Pathways</b>	Neuroactive ligand-receptor interaction
<b>Molecular Weight</b>	The human full length GIPR-Strep protein has a MW of 53.2 kDa
<b>Delivery</b>	6~8weeks
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Sterility</b>	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	A G-protein coupled receptor for gastric inhibitory polypeptide (GIP), which was originally identified as an activity in gut extracts that inhibited gastric acid secretion and gastrin release, but subsequently was demonstrated to stimulate insulin release in the presence of elevated glucose. Mice lacking this gene exhibit higher blood glucose levels with impaired initial insulin response after oral glucose load. Defect in this gene thus may contribute to the pathogenesis of diabetes.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

