

## PRODUCT INFORMATION

<b>Target</b>	GITR Ligand
<b>Synonyms</b>	TNFSF18;AITRL;TL6;hGITRL;GITR Ligand
<b>Description</b>	Recombinant human GITR Ligand protein with N-terminal mouse Fc and C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9UNG2
<b>Expression Host</b>	HEK293
<b>Tag</b>	N-Mouse Fc and C-6×His Tag
<b>Molecular Characterization</b>	mFc(Pro99-Lys330) GITR Ligand(Gln50-Ser177) 6×His
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 46-53 kDa after removal of the signal peptide.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage&amp;Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptor TNFRSF18/AITR/GITR. It has been shown to modulate T lymphocyte survival in peripheral tissues. This cytokine is also found to be expressed in endothelial cells, and is thought to be important for interaction between T lymphocytes and endothelial cells.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated







Figure 1. Human GITR Ligand Protein, mFc-His Tag on SDS-PAGE under reducing condition.

Figure 2. Flow cytometry analysis with 1  $\mu$ g/mL Human GITR Ligand Protein, mFc-His tag (PME100020) on HEK293 cells transfected with human GITR (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).