

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag Tag |
| Expression Host | HEK293 |
| Target | GLP1R |
| Synonyms | GLP-1; GLP-1-R; GLP-1R |
| Description | Human GLP1R full length protein-synthetic nanodisc |
| Uniprot ID | P43220 |
| Protein Families | Druggable Genome, ES Cell Differentiation/IPS, GPCR, Transmembrane |
| Protein Pathways | Neuroactive ligand-receptor interaction |
| Molecular Weight | The human full length GLP1R protein has a MW of 53.0 kDa |
| Delivery | In Stock |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Sterility | Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use. |
| Storage&Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | A 7-transmembrane protein that functions as a receptor for glucagon-like peptide 1 (GLP-1) hormone, which stimulates glucose-induced insulin secretion. This receptor, which functions at the cell surface, becomes internalized in response to GLP-1 and GLP-1 analogs, and it plays an important role in the signaling cascades leading to insulin secretion. It also displays neuroprotective effects in animal models. Polymorphisms in this gene are associated with diabetes. The protein is an important drug target for the treatment of type 2 diabetes and stroke. Alternative splicing of this gene results in multiple transcript variants. |
| Usage | Research use only |
| Conjugate | Unconjugated |



ELISA assay to evaluate GLP1R-Nanodisc 0.2 μ g Human GLP1R-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag GLP1R-Nanodisc (0.2 μ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GLP1R-Nanodisc is 3.549ng/ml.



Figure2. Human GLP1R-Nanodisc, Flag Tag on SDS-PAGE

