

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	GPR17
Synonyms	Uracil nucleotide/cysteinyl leukotriene receptor; UDP/CysLT receptor; G-protein coupled receptor 17; P2Y-like receptor; R12
Description	Human GPR17-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q13304
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,GPCRDB Other,
Molecular Weight	The human full length GPR17-Strep protein has a MW of 41 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Sterility	Products are supplied non-sterile. For cell culture applications, dilute in appropriate medium and sterile-filter (0.22 µm) prior to use.
Background	Dual specificity receptor for uracil nucleotides and cysteinyl leukotrienes (CysLTs). Signals through G(i) and inhibition of adenylyl cyclase. May mediate brain damage by nucleotides and CysLTs following ischemia.[UniProtKB/Swiss-Prot Function]
Usage	Research use only
Conjugate	Unconjugated

