Cat. No. FLP100031

Delivery

Background



PRODUCT INFORMATION

Target GPR75

Synonyms GPRchr2; WI31133

Human GPR75 full length protein-synthetic **Description**

nanodisc In Stock

Uniprot ID 095800 **Expression Host HEK293**

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways

The human full length GPR75 protein has a MW of **Molecular Weight**

59.4 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

G protein-coupled receptor that is activated by the chemokine CCL5/RANTES. Probably coupled to heterotrimeric Gq proteins, it stimulates inositol trisphosphate production and calcium mobilization upon activation. Together with

CCL5/RANTES, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. CCL5/RANTES may also regulate insulin secretion by page 25th is left cells through activation of this by pancreatic islet cells through activation of this

réceptor.

Usage Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com





ELISA assay to evaluate GPR75-Nanodisc 0.2µg Human GPR75-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GPR75-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-GPR75 monoclonal antibody (DMC100368) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-GPR75 monoclonal antibody binding with GPR75-Nanodisc is 303.6ng/ml.




Figure 2. WB analysis of Human GPR75-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

Email: info@dimabio.com Website: www.dimabio.com

