**Delivery** 



## **PRODUCT INFORMATION**

**Target** C5AR2

C5L2; GPF77; GPR77 **Synonyms** 

Human GPR77 full length protein-synthetic **Description** 

nanodisc 3-4 weeks

**Uniprot ID** Q9P296 **Expression Host HEK293** 

**Protein Families** Druggable Genome, GPCR, Transmembrane

**Protein Pathways** 

The human full length C5AR2 protein has a MW of **Molecular Weight** 

36.1 kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution for specific instructions. Do not use solvents with

pH lower than 6.5 in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

Storage & Shipping at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a G-protein coupled receptor 1 family member involved in the complement system of the innate immune response. Unlike classical G-protein coupled receptors, the encoded protein does not associate with

intracellular G-proteins. It may instead modulate **Background** signal transduction through the beta-arrestin

pathway, and may alternatively act as a decoy receptor. This gene may be involved in coronary artery disease and in the pathogenesis of sepsis. Alternative splicing results in multiple transcript

> Email: info@dimabio.com Website: www.dimabio.com

variants. [provided by RefSeq, Nov 2012]

**Usage** Research use only

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





## ELISA assay to evaluate GPR77-Nanodisc 0.2µg Human GPR77-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag GPR77-Nanodisc ( $0.2\mu g/per$  well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GPR77-Nanodisc is 3.415 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com



Figure 2. Human GPR77-Nanodisc, Flag Tag on SDS-PAGE

