

**PRODUCT INFORMATION**

|   |  |
|---|--|
| <b>Tag</b>                              | C-Flag Tag   |
| <b>Target</b>                           | GPR84  |
| <b>Synonyms</b>                         | EX33; GPCR4  |
| <b>Description</b>                      | Human GPR84 full length protein-synthetic nanodisc   |
| <b>Delivery</b>                         | In Stock   |
| <b>Uniprot ID</b>                       | Q9NQ55   |
| <b>Expression Host</b>                  | HEK293   |
| <b>Protein Families</b>                 | Druggable Genome, GPCR, Transmembrane  |
| <b>Protein Pathways</b>                 | N/A  |
| <b>Molecular Weight</b>                 | The human full length GPR84 protein has a MW of 43.7 kDa   |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.   |
| <b>Storage&amp;Shipping</b>             | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.  |
| <b>Background</b>                       | Receptor for medium-chain free fatty acid (FFA) with carbon chain lengths of C9 to C14. Capric acid (C10:0), undecanoic acid (C11:0) and lauric acid (C12:0) are the most potent agonists. Not activated by short-chain and long-chain saturated and unsaturated FFAs. Activation by medium-chain free fatty acid is coupled to a pertussis toxin sensitive G(i/o) protein pathway. May have important roles in processes from fatty acid metabolism to regulation of the immune system. |
| <b>Usage</b>                            | Research use only  |
| <b>Conjugate</b>                        | Unconjugated   |



### ELISA assay to evaluate GPR84-Nanodisc 0.2 $\mu$ g Human GPR84-Nanodisc per well




Figure1. Elisa plates were pre-coated with Flag Tag GPR84-Nanodisc (0.2 $\mu$ g/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GPR84-Nanodisc is 2.893ng/ml.



Figure2. WB analysis of Human GPR84-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

